

NAM
Acorn House
314-320 Gray's Inn Road
London
WC1X 8DP
Phone: 020 7837 6988
Website: www.aidsmap.com
Email: info@nam.org.uk
Charity number: 1011220

Thanks to all of the professionals and people
with HIV who have helped us to develop
this leaflet.

nam aidsmap

The basics

Undetectable viral load and HIV transmission

Notes

Find out more

For more information on this topic

- Read our 'Undetectable viral load' leaflet.
- Read 'Viral load and transmission' on our website.
- Read our 'HIV & sex' booklet.

For more information about HIV

- Visit our website at www.aidsmap.com.
- Speak to an adviser at THT Direct (phone: 0808 802 1221).

For information in other languages


- Visit www.aidsmap.com/translations.

Has this leaflet been useful to you?

Please let us know what you think. Your feedback helps us to improve the services we offer.

You can contact us to find out more about the scientific research and information we have used to produce this leaflet.

We recommend that you discuss the information in this leaflet with a doctor or other health worker.




First edition 2019
Due for review in 2022

Copyright ©NAM


If you need extra copies of this leaflet, or would like to translate, adapt or reproduce it, please contact us.

Effective HIV treatment stops HIV from being passed on during sex.

HIV treatment reduces the amount of HIV in the body. If you have an 'undetectable viral load', this means that there is not enough HIV to pass on to anybody else.



If you are taking HIV treatment and have an undetectable viral load, you will **not** pass HIV on during sex.




You can rely on an undetectable viral load if

- your viral load has been undetectable for at least six months; and
- you don't miss doses of your treatment.



We know this from scientific studies of couples where one partner had HIV and the other did not.

- The couples had sex without a condom **130,000 times**.
- There were **no cases** of HIV being passed on.



You do not need to use condoms to prevent HIV transmission when your viral load is undetectable. But condoms will help to prevent other sexually transmitted infections.



Important points

- **If you have had an undetectable viral load for at least six months and you continue to take your treatment as prescribed, there is no risk of passing HIV on during sex.**
- **This is why people say 'undetectable = untransmittable' (U=U).**